REFERENCES

- Adriana L.M.A., Jayme A.A.S., Elfriede M.B., (2002), Antiulcer activity of Sapindus saponaria L. in the rat, Journal of Ethnopharmacology, 82: 41-44.
- Agrawal R., Garg H.K., Garg U., Singh S.K., (2010), Anti-ulcer activity of Smithia conferta in various animal, Journal of Saudi Chemical Society, 14: 307–310.
- Ahmed N., (2005), Advanced glycation endproducts-role in pathology of diabetic complications. Diabetes Research and Clinical Practice, 67; 3–21.
- Alam S., Mohammed A., Syed M.B.A., Satya P.V., (2009), Antiulcer activity of methanolic extract of *Momordica charantia* L. in rats, Journal of Ethnopharmacology, 123: 464–469.
- Alimia H., Najla H., Zouhour B., Mbarka H., Mohsen S., Lazhar Z., Khémais B.R., (2010), Antioxidant and antiulcerogenic activities of Opuntia ficus indica f. inermis root extract in rats, Phytomedicine, 17: 1120–1126.
- Althunibat O.Y., Ahmed H.A., Khaled T., Khaled M. K., Ridzwan B.H., Haitham N.Q., (2010), Protective role of Punica granatum L. peel extract against oxidative damage in experimental diabetic rats, Process Biochemistry, 45: 581–585.
- Arawwawalaa L.D.A.M., Thabrewb M.I., Arambewela L.S.R.,, (2010), Gastroprotective activity of Trichosanthes cucumerina in rats, Journal of Ethnopharmacology, 127: 750–754
- Azadbakhta M., Safapour S., Ahmadi A., Maryam G., Shokrzadeh M., (2010), Anti-diabetic effects of aqueous fruits extract of *Diospyros lotus L*. on streptozotocin-induced diabetic rats and the possible morphologic changes in the liver, kidney and heart, Journal of Pharmacognosy and Phytotherapy, Vol. 2(2): 10-16
- Babincová M., Kristína S., Paul S., (2008), Antiulcer activity of water extract of Scoparia dulcis, Fitoterapia, 79: 587–588.
- Barbastefano V., Cola M., Luiz-Ferreira A., Farias-Silva E., Hiruma-Lima C.A., Rinaldo D., Wagner V., Alba R.M., (2007), Vernonia polyanthes as a new source of antiulcer drugs, Fitoterapia, 78: 545–551.

- Chandramohan G., Al-Numair K.S., Sridevi M., Pugalendi K.V., (2008), A novel compound from Casearia esculenta (Roxb.) root and its effect on carbohydrate metabolism in streptozotocin-diabetic rats, European Journal of Pharmacology, 590: 437–443.
- Chatterjee A and Pakrashi S.C., (1992), The treatise of Indian medicinal plants, Pp 180.
- Chaturvedi A., Bhawani G., Agarwal P.K., Goel S, Singh A., Goel R.K., (2009), Antidiabetic and antiulcer effects of extract of *Eugenia jambolana* seed in mild diabetic rats: Study on gastric mucosal offensive Acid-pepsin secretion, Indian J Physiol Pharmacol, 53 (2): 137–146.
- DeAndrade S. F., Marivane L., Eros C., Vânia F. N., Valdir C. F., Rivaldo N., (2007), Evaluation of the antiulcerogenic activity of Maytenus robusta (Celastraceae) in different experimental ulcer models. Journal of Ethnopharmacology, 113; 252–257.
- Falca˜ o H.S., Mariath I.R., Diniz M.F.F.M., Batista L.M., and Barbosa-Filho J.M., (2008), Plants of the American continent with antiulcer activity, Phytomedicine, 15: 132–146.
- Ganapathy S., Ponnusamy P., Gandhipuram P., Senthil K., Thatchinamoorthi R., (2009), Modulatory effect of S-allylcysteine on glucose metabolism in streptozotocin induced diabetic rats. Journal of functional foods, 1; 336-340.
- Gilani A., Bashir S., Janbaz K.H., Shah A.J.,(2005), Presence of cholinergic and calcium channel blocking activities explains the traditional use of *Hibiscus rosasinensis* in constipation and diarrhoea, Journal of Ethnopharmacology, 102: 289–294.
- Goel R.K. and Sairam K. (2002), Antiulcer drugs from indigenous sources with emphasis on musa sapientum, tamrabhasma, asparagus racemosus and Zingiber officinale. Indian Journal of pharmacology, 34; 100-110.
- Goulart Y.C.F., Vania R.S., Simoni O., Martins J.V.C., Fernanda O., Cortez D.A., Elisabeth A.A., (2005), Evaluation of Gastric Anti-ulcer Activity in a Hydroethanolic Extract from *Kielmeyera coriacea.* Brazilian Archives of Biology and Technology, Vol.48; 211-216,

- Grover J.K.,, Yadav S.P., Vats V., (2003), Effect of feeding *Murraya koeingii* and *Brassica juncea* diet kidney functions and glucose levels in streptozotocin diabetic mice, Journal of Ethnopharmacology, 85: 1–5.
- Guaraldo L., Sertie`J. A.A., Bacchi E.M., (2001), Antiulcer action of the hydroalcoholic extract and fractions of *Davilla rugosa* Poiret in the rat, Journal of Ethnopharmacology, 76: 191–195.
- Hiruma-Lima C.A., (2006), Antiulcerogenic activity of Alchornea castaneaefolia: Effects on somatostatin, gastrin and prostaglandin. Journal of Ethnopharmacology, 104; 215–224.
- Jain S.M., Parmar N.S., Santani D.D., (1994), Gastric Antiulcer Activity of Calcium Channel Blockers in Rats, Indian Journal of Pharmacology, 26: 29 – 34.
- Jamal A., Javed K., Aslama M., Jafri M.A., (2006), Gastroprotective effect of cardamom, Elettaria cardamomum Maton. fruits in rats, Journal of Ethnopharmacology, 103: 149–153.
- Joseph J. M., Kandhasamy S., Sellamuthu M., (2010), Protective effects of methanolic extract of Hedyotis puberula (G. Don) R. Br. Ex Arn. against experimentally induced gastric ulcers in rat, Journal of Ethnopharmacology, 131: 216–219.
- Kanai S., (2007), Decreased hydrogen–potassium-activated ATPase (H+–K+-ATPase) expression and gastric acid secretory capacity in aged mice. Archives of Gerontology and Geriatrics, 45; 243–252.
- Konan N. A., Elfriede M.B., (2007), Antiulcerogenic effect and acute toxicity of a hydroethanolic extract from the cashew (Anacardium occidentale L.) leaves, Journal of Ethnopharmacology, 112: 237–242.
- Lee J.S., (2006), Effects of soy protein and genistein on blood glucose, antioxidant enzyme activities, and lipid profile in streptozotocin-induced diabetic rats. Life Sciences, 79; 1578–1584
- Luiz-Ferreira A., (2008), Should Anacardium humile St. Hil be used as an antiulcer agent? A scientific approach to the traditional knowledge. Fitoterapia, 79; 207–209.
- Maity B and Chattopadhyay S. (2008), Natural Antiulcerogenic Agents: An Overview. Current bioactive compounds, 4; 225-244.

- Manonmani S., (1995), Biochemical studies on the antiulcerogenic activity of Cauvery 100, an ayurvedic formulation in experimental ulcers. Indian Journal of Pharmacology, 27: 101 – 105.
- Mes´ıa-Vela S., (2007), In vivo inhibition of gastric acid secretion by the aqueous extract of Scoparia dulcis L. in rodents. Journal of Ethnopharmacology, 111; 403–408.
- Nirmala A., Saroja S., Vasanthi H.R., Lalitha G., (2009), Hypoglycemic effect of *Basella rubra* in streptozotocin – induced diabetic albino rats. Journal of Pharmacognosy and Phytotherapy Vol. 1 (2) pp. 025-030.
- Pandit R., Phadke A., Jagtap A., (2010), Antidiabetic effect of Ficus religiosa extract in streptozotocin-induced diabetic rats. Journal of Ethnopharmacology, 128; 462–466.
- Pascual M.E., (2001), Antiulcerogenic activity of *Lippia alba* (Mill.) N. E. Brown (Verbenaceae). Il Farmaco, 56; 501–504.
- Perera L.M.S., Dayamı´ R., Go´mez B.C., (2001), Gastric antiulcer effect of Rhizophora mangle L. Journal of Ethnopharmacology, 77: 1–3
- Rai V., Iyer U., Mani U.V., (1997), Effect of Tulasi (Ocimum sanctum) leaf powder upplementation on blood sugar levels, serum lipids and tissue lipids in diabetic rats. Plant Foods Hum Nutr, 50; 9-16.
- Ramesh B., Periyasamy V., Kodukkur V.P., (2007), Protective effect of Umbelliferone on membranous fatty acid composition in streptozotocin-induced diabetic rats, European Journal of Pharmacology, 566: 231–239.
- Rao Ch. V., Ojha S.K., Radhakrishnan K., Govindarajan R., Rastogi S., (2004), Antiulcer activity of Utleria salicifolia rhizome extract, Journal of Ethnopharmacology, 91: 243–249.
- Shamburek, R.D; Schubert, M.L. (1992). Control of gastric acid secretion. Histamine H_2 – receptor antagonists and H + K(+) – ATPase inhibitors. Gastroenterolol. *Clin. North Am*. 21:527-550
- Shine V.J., Latha P.G., Shyamal S., Suja S.R., Anuja G.I., Sini S., Pradeep S., Rajasekharan S., (2009), Gastric antisecretory and antiulcer activities of *Cyclea peltata* (Lam.) Hook. f. & Thoms. in rats, Journal of Ethnopharmacology, 125: 350–355

- Sohn E., (2010), Extract of the aerial parts of Aster koraiensis reduced development of diabetic nephropathy via anti-apoptosis of podocytes in streptozotocin-induced diabetic rats. Biochemical and Biophysical Research Communications, 391; 733–738.
- Souccar C, Cysneiros R.M., Tanae M.M., Torres L.M.B., Lima-Landman M.T.R., Lapa A.J., (2008), Inhibition of gastric acid secretion by a standardized aqueous extract of Cecropia glaziovii Sneth and underlying mechanism, Phytomedicine, 15: 462–469
- Souza M.C., (2009), In vitro and in vivo anti-Helicobacter pylori activity of Calophyllum brasiliense Camb. Journal of Ethnopharmacology, 123; 452–458.
- Suffredini I. B., Elfriede M. B., Jayme A.A.A. S., (1999), Antiulcer action of *Microgramma squamulosa* (Kaulf.) Sota Journal of Ethnopharmacology, 65:217–223.
- Wahida B., Abderrahman B., Nabil C., (2007), Antiulcerogenic activity of *Zizyphus lotus* (L.) extracts, Journal of Ethnopharmacology, 112: 228–231.
- Wang J., Zhu L., Zou K., Cheng F., Dan F., Guo Z., Cai Z., Yang J., (2009), The anti-ulcer activities of bisabolangelone from *Angelica polymorpha.* Journal of Ethnopharmacology, 123; 343–346.